


# CASE STUDY

**PHAM MINH THONG MD**  
**PROFESSOR OF RADIOLOGY**  
**Bachmai University Hospital**  
**Hanoi, Vietnam**

# Case No2

- F, 56 YO, headache
- Basilar tip aneurysm
- Non-rupture
- Wide-neck
- Round shape




# Treatment

- Surgery?
- Coiling direct?
- Remodelling technique?
- Stent assisted coiling?

# Treatment

- Surgery?
- Coiling direct?
- Remodelling technique?
- Stent assisted coiling?




- Stent Lvis 2,5x18 in the Basilar and right P1
- Jailed microcatherter technique




- Ruptured during deploying the second coils

What should we do?



- Push balloon hyperform inside the stent to inflate, continue coiling



**Control:** stop  
bleeding and  
Aneurysm  
occluded

- Wake-up: Mild headache.
- CT scanner: Mild SAH
- Discharge hospital after 5 days:  
mRS 0

# Teaching points

- Non rupture Wide –neck aneurysm in the tip of basilar trunk: stent + coiling
- Rupture of Aneurysm during the procedure: Neutralization of Heparin and balloon assisted necessary to stop bleeding
- Continue coiling quickly and maintain Heparin after total occlusion of the Aneurysm